

## Clinical Study

# A study of retroperitoneal core needle biopsy in the diagnosis of suspected malignant renal tumors

 Priya Singh<sup>1</sup>, Yogesh Kumar Sarin<sup>1</sup>, Anita Nangia<sup>2</sup>

 Departments of Pediatric Surgery<sup>1</sup> and Pathology<sup>2</sup>, Lady Hardinge Medical College, New Delhi-110001.

**Keywords**

Clear cell sarcoma of kidney  
 Core needle biopsy  
 Kidney tumors in children  
 Malignant rhabdoid tumor  
 Non-Wilms renal tumor  
 Renal tumor  
 Wilms tumor

**Abbreviations**

**CCSK** - Clear cell sarcoma of kidney  
**CECT** - Contrast enhanced computed tomography  
**CNB** - Core needle biopsy  
**COG** - Children's oncology group  
**HPE** - Histopathological examination  
**MRTK** - Malignant rhabdoid tumor of kidney  
**NACT** - Neo-adjuvant chemotherapy  
**NWRT** - Non-Wilms renal tumors  
**NWTSG** - National Wilms tumor study group  
**RTGS** - Renal tumor study group  
**SIOP** - Societe Internationale D'Oncologie Pediatric  
**USG** - Ultrasonography  
**WT** - Wilms tumor

**Abstract**

**Background:** Differentiation of Wilms tumor (WT) from non-Wilms renal tumors (NWRT) is essential as the management philosophies of these two groups of tumors are very different, especially if we believe in the SIOP protocol of administering neo-adjuvant chemotherapy (NACT) in patients with WT. This study examines the role of the preoperative retroperitoneal core needle biopsy (CNB) in the differential diagnosis of suspected malignant renal tumors in children

**Material and methods:** CNB was done in 20 patients (23 renal biopsies as 3 patients had bilateral tumors) with suspected malignant renal tumors. Those diagnosed as WT on CNB received NACT followed by nephrectomy and adjuvant therapy. We studied different subtypes of WT, chemotherapy-induced changes in the resected specimen and the concordance between the histopathology examination (HPE) of pre-operative CNB with that of the excised surgical specimen.

**Results:** Out of the 23 renal biopsies, 21 were diagnosed as WT while the other 2 as malignant rhabdoid tumors of kidney (MRTK). One patient developed abdominal distension and respiratory distress after CNB. On HPE of the excised surgical specimen of those who received NACT for WT (diagnosed on CNB), 19 samples were confirmed as WT, while 2 were diagnosed as clear cell sarcoma of the kidney (CCSK). Of the 2 tumors that were diagnosed as MRTK on CNB, 1 was confirmed as MRTK and the other was diagnosed as CCSK on HPE of the excised specimen. The concordance between the HPE of CNB and the excised surgical specimen was 91%.

**Conclusion:** Pre-therapy CNB is a sensitive tool to differentiate WT from NWRT. The sensitivity, specificity, positive predictive value and negative predictive value of CNB in the diagnosis of renal tumors are 89%, 50%, 89% and 100% respectively.

## INTRODUCTION

Childhood cancers represent a significant health concern worldwide, with malignant renal tumors accounting for approximately 7% of all childhood cancers. Among these, Wilms Tumor (WT) is one of the most common, accounting for about 85% of all pediatric malignant renal tumors.<sup>(1)</sup> Accurate diagnosis of renal tumors is crucial for selecting the most appropriate treatment and ensuring optimal outcomes for patients. It is not always possible to differentiate between different types of renal tumors by clinical examination and imaging alone.<sup>(2)</sup> Saula *et al.* noted approximately 13.6% of cases were NWRT in the developing countries, highlighting the importance of preoperative CNB in guiding treatment decisions.<sup>(3)</sup> Moreover, NWRT are far more commonly seen in Southeast Asian countries including India as compared to the Western world.<sup>(3)</sup> WT is treated according to the guidelines suggested by the National Wilms tumor study group (NWTSG) - recently renamed as Children's Oncology Group (COG) - or the International Society of Pediatric Oncology (SIOP).<sup>(4)</sup> NWTSG/COG believes in up-front surgery followed by adjunct therapy, while SIOP believes in neo-adjuvant chemotherapy (NACT) followed by surgery and then adjuvant therapy.


During SIOP trials, it was noted that 5-10% of tumors initially suspected to be WT turned out to be either benign renal tumors or NWRT that do not respond to NACT.<sup>(5)</sup> COG approach has the advantage of studying tumor histology unaltered by prior chemotherapy and hence it makes subtyping and staging of the tumor easier. Even in COG, NACT is administered to inoperable WT, WT in a solitary kidney, synchronous bilateral WT, tumor thrombus in inferior vena cava extending above the level of the hepatic veins and the tumor involving contiguous vital structure.

In India, pre-treatment CNB is not a popular practice, unlike in the United Kingdom where the Children's Cancer Study Group (UKCCSG) used to advocate preoperative CNB through the retro-

peritoneal route for all suspected WT to ascertain the histopathological diagnosis before administering neo-adjuvant chemotherapy.<sup>(5)</sup> As the management philosophy of WT and NWRT is very different, we intended to assess the role of the pre-therapy retroperitoneal CNB in the differential diagnosis of pediatric renal.

## MATERIALS AND METHODS

After obtaining the approval of the institute ethical committee (LHMC/IEC/2021/03/88), a prospective descriptive study of 20 patients with suspected malignant renal tumors was done during the period of July 2021 to November 2022. Patients with recurrent disease and tumor rupture or hemorrhage necessitating emergency surgery were excluded. All the patients had routine biochemical investigations, ultrasonography (USG) abdomen with Doppler and contrast enhanced computed tomographic scan (CECT) of the chest and abdomen. (Fig. 1)



**Fig 1. Left renal Tumor**  
Contrast Enhanced Computed Tomography (CECT) showing the left renal mass in axial (upper panel) and coronal (lower panel) section.



**Fig. 2: USG guided core-needle biopsy procedure.**

All but one patient underwent USG-guided CNB through the retro-peritoneal route after administering adequate sedation and local anesthesia; one patient erroneously underwent transperitoneal CNB.(Fig.2) At least 2-3 cores were taken and sent for HPE. Adverse events, either during or after the procedure, were recorded. HPE was reported within approximately 5-6 days. For patients with bilateral renal tumors, both tumors were biopsied as discordant pathologies are known to occur. Those who were diagnosed as WT were given NACT as per the Umbrella protocol of SIOP-RTSG<sup>(18)</sup> and those diagnosed as MRTK underwent upfront surgery. In CNB diagnosed WT, a re-evaluation was done by CECT of the abdomen before nephrectomy to assess the reduction in tumor volume due to NACT.

Written consent for surgery was obtained from caregivers. All patients underwent tumor excision by a single senior surgeon (YKS). Sixteen patients underwent total nephro-ureterectomy; these include 2 children diagnosed to have MRTK on CNB. Four patients underwent nephron-sparing surgery; 3 of them had bilateral WT. Intra-operative and post-operative adverse events, if any, were recorded. The excised surgical specimen was sent for HPE (Fig. 3) to study tumor staging and risk-stratification. Concordance between the pre-therapy CNB report and that of post-nephrec-



**Fig. 4: Left nephro-ureterectomy specimen**

tomy HPE was recorded. NACT-induced changes in histopathology were noted. In the case of bilateral tumors, adjuvant therapy was based on the tumor having the higher risk stratification.

## RESULTS

CNB was performed for 23 renal tumors in 20 patients; 3 had bilateral renal tumors. In one patient, the CNB was done transperitoneally by mistake. She developed abdominal distension, respiratory distress and a fall in hemoglobin due to intra-abdominal bleeding. She was upstaged to stage-3 and was treated accordingly. CNB-related hemorrhage, tumor rupture or tumor seeding was not noted in any other patient.

Eighteen patients including 3 having bilateral WT (21 renal CNB) were diagnosed to have WT. They received NACT as per SIOP protocol. The other 2 patients diagnosed as MRTK on CNB underwent upfront nephrectomy.(Table 1). More than half of the WT (57.1%) patients were reported to have biphasic WT on CNB.

The interpretation of monophasic WT with blastemal component alone (n=2) on CNB was easy to interpret. However, interpreting monophasic WT with stromal component (n=2) needed considerable expertise. In 19% (4/21) rhabdomyoblastic changes were seen.

Table 1: Histopathological diagnosis on core-needle biopsy

Renal tumors	n (%)	Blastemal	Epithelial	Stromal	RMB
WT	Monophasic 4 (19%)	2	-	2	-
	Biphasic 12 (57.1%)	8	9	7	3
	Triphasic 5 (23.8%)	5	5	5	1
MRTK	2 (8.6%)	-	-	-	-

MRTK - Malignant rhabdoid tumor of kidney, RMB - Rhabdo-myoblastic change; WT - Wilms tumor

Table 2: Histopathological diagnosis of the nephrectomy specimens

Tumor Histology with risk stratification	n (%)	Blastemal	Epithelial	Stromal	RMB
WT (n=19; 82.6%)					
IR-17					
Mixed type - 9	Monophasic 1 (5.2%)	-	1	-	-
Stromal type - 4	Biphasic 5 (26.3%)	-	5	5	2
Regressive type - 3					
Epithelial type - 1	Triphasic 13 (68.4%)	13	13	13	3
HR (Blastemal type) - 2					
CCSK (HR)	3 (13%)	-	-	-	-
MRTK (HR)	1 (4.3%)	-	-	-	-

CCSK- Clear cell sarcoma of kidney; HR- High risk; IR- Intermediate risk; MRTK- Malignant rhabdoid tumor of kidney; RMB - Rhabdo-myoblastic change; WT-Wilms tumor.

None of the WT showed anaplasia. The mean reduction of tumour volume after NACT was 41.8  $\pm$  2%. No change in the tumor volume after NACT was noted in 13/21 (61.9%) tumors. None of the bilateral tumors (n=6) or the 2 tumors that later turned out to be CCSK showed any change in tumor volume. Only 3 patients with unilateral WT on CNB showed considerable (>50%) reduction in tumor volume following NACT. Heterologous elements like cartilage and bone were found in 2 specimens, while rhabdo-myoblastic changes were seen in 5 specimens.

On histopathology of the excised specimens, 17 (73.9%) patients had intermediate-risk tumors and 6 (26%) had high-risk tumors as per working SIOP classification; none was of 'low-risk' category (completely necrotic).

In 2 out of the 21 WT and one out of the 2 MRTK diagnosed on CNB, the diagnosis was revised as CCSK on the final histology of nephrectomy specimen.(Table 2) The concordance between CNB and histology of resected specimen was 91%. The sensitivity, specificity, positive predictive value and negative predictive value of CNB in the

diagnosis of pediatric renal tumors were 89%, 50%, 89% and 100% respectively.

About three-fourths of the 19 excised surgical specimens had <33% viable tumor cells reported as the blastemal component, thus proving that the blastemal component is chemo-sensitive. Only 2 specimens had shown blastemal component of >66% thus designating it as a high-risk tumor with a poor prognosis.

## DISCUSSION

In the field of pediatric oncology, obtaining an accurate diagnosis in the shortest period of time with minimal morbidity is of utmost importance.<sup>(17)</sup> Of all childhood cancers, malignant renal tumors comprise 7% of all cancers. Among them, WT is the most common renal tumor which is commonly presented in the age group from 1-4 years.<sup>(1,2,6)</sup> The differentiation between WT and NWRT can be challenging as they share many radiological features. Therefore, CNB is necessary for accurate diagnosis. UKCCSG WT Study 3 adopted pre-NACT CNB for histological diagnosis to address these issues [13].

The diagnosis of WT and other renal tumors is typically made by imaging studies, including USG (including Doppler), computed tomography and magnetic resonance imaging. These studies can provide detailed information about the size, location and characteristics of the tumor, which are invaluable in therapeutic decision making. In some cases, a CNB may be performed to confirm the diagnosis, but this is not always necessary, especially when the imaging studies are highly suggestive of a specific type of tumor.<sup>(5)</sup> According to Umbrella protocol, CNB is now considered only in cases with unusual clinical presentation like age >6 years, urinary infection or septicemia, psoas infiltration, pulmonary metastasis in <2 years, and extra-hepatic or extra-pulmonary metastasis, unusual radiological findings like numerous calcifications, voluminous lymphadenopathy, non visualization of renal parenchyma, extra-renal

projections and biological findings like hypercalcemia or LDH level >4 x ULN (upper limit of normal).<sup>(18)</sup> It is our institutional policy to do CNB in every patient suspected to have a malignant renal tumor.

Skoldenberg's study showed that the sensitivity of CNB was 76%.<sup>(14)</sup> Taskinen *et al.* reported a sensitivity of 100% for CNB in diagnosing WT; however, only two out of five NWRT cases in their series were diagnosed with CNB.<sup>(4)</sup> Jackson *et al.* showed that the sensitivity and specificity of CNB were 86% and 99.6%, respectively. In 25 out of 518 cases, CNB changed the management, thereby demonstrating the value of this procedure.<sup>(15)</sup> Mitchell *et al.*<sup>(16)</sup> reported that approximately 12% of cases were diagnosed as NWRT due to CNB and were treated accordingly, even though they had features suggestive of WT in radiological studies.

## CONCLUSION

We conclude that in our experience CNB is an effective tool in diagnosing WT and differentiating it from NWRT. Inappropriate NACT has been avoided in 2 patients who were diagnosed with NWRT on CNB. In 3 bilateral WT, NACT prescribed by SIOP Umbrella protocol didn't reduce any tumour volume, which indicates that we should consider administering 3-drug NACT as suggested in COG.

## REFERENCES

- [1] Lee JS, Sanchez TR, Wootton-Gorges S. Malignant renal tumors in children. *J Kidney Cancer VHL*. 2015 May 10; 2 (3): 84-89.
- [2] Chung EM, Graeber AR, Conran RM. Renal Tumors of Childhood: Radiologic-Pathologic Correlation Part 1. The 1st Decade: From the Radiologic Pathology Archives. *Radiographics*. 2016 Mar-Apr; 36(2): 499-522.
- [3] Saula PW, Hadley GP. Pediatric non-Wilms renal tumors: a third world experience. *World J Surg*. 2012 Mar; 36 (3): 565-72.
- [4] Taskinen S, Lohi J, Koskenvuo M, Taskinen M. Evaluation of effect of preoperative chemotherapy on Wilms tumor histopathology. *J Pediatr Surg*. 2018 Aug; 53(8): 1611-1614.
- [5] Vujanić GM, Kelsey A, Mitchell C, Shannon RS, Gornall P. The role of biopsy in the diagnosis of renal tumors of

childhood: Results of the UKCCSG Wilms tumor study 3. *Med Pediatr Oncol.* 2003 Jan; 40 (1): 18-22.

[6] Charles AK, Vujanić GM, Berry PJ. Renal tumours of childhood. *Histopathology.* 1998 Apr; 32(4): 293-309.

[7] Lemerle J, Voute PA, Tournade MF, Rodary C, Delemarre JF, Sarrazin D, Burgers JM, Sandstedt B, Mildenberger H, Carli M, Jereb B, Moorman-Voestermans CGM. Effectiveness of preoperative chemotherapy in Wilms tumor: results of an International Society of Paediatric Oncology (SIOP) clinical trial. *J Clin Oncol.* 1983 Oct; 1(10): 604-9.

[8] Saarinen UM, Wikström S, Koskimies O, Sariola H. Percutaneous needle biopsy preceding preoperative chemotherapy in the management of massive renal tumors in children. *J Clin Oncol.* 1991 Mar; 9(3): 406-15.

[9] Sarin YK, Sinha S, Khurana N. Chemotherapy-induced changes in Wilms tumor - our experience. *Pediatr Blood Cancer* 2014; 61: S408.

[10] Dykes EH, Marwaha RK, Dicks-Mireaux C, Sams V, Risdon RA, Duffy PG, Ransley PG, Pritchard J. Risks and benefits of percutaneous biopsy and primary chemotherapy in advanced Wilms tumour. *J Pediatr Surg.* 1991 May; 26 (5): 610-2.

[11] Greenberg M, Burnweit C, Filler R, Weitzman S, Sohl H, Chan H, Jenkin D, Dougherty M, Berry M, Balcolm A, McLorie G. Preoperative chemotherapy for children with Wilms tumor. *J Pediatr Surg.* 1991 Aug; 26(8): 949-56.

[12] Zoeller G, Pekrun A, Lakomek M, Ringert RH. Wilms tumor: the problem of diagnostic accuracy in children undergoing preoperative chemotherapy without histological tumor verification. *J Urol.* 1994 Jan; 151(1): 169-71.

[13] United Kingdom Children's Cancer Study Group. 1992. Wilms tumour study 9101 (UKW3) protocol. {www.ukccs.org} (Accessed 27 February 2024)

[14] Sköldenberg EG, Jakobson A, Elvin A, Sandstedt B, Läckgren G, Christofferson RH. Pretreatment, ultrasound guided cutting needle biopsies in childhood renal tumors. *Med Pediatr Oncol.* 1999 Apr; 32(4): 283-8.

[15] Jackson TJ, Williams RD, Brok J, Chowdhury T, Ronghe M, Powis M, Pritchard-Jones K, Vujanić GM; Children's Cancer and Leukaemia Group (CCLG) Renal Tumours Group. The diagnostic accuracy and clinical utility of pediatric renal tumor biopsy: Report of the UK experience in the SIOP UK WT 2001 trial. *Pediatr Blood Cancer.* 2019 Jun; 66 (6): e27627.

[16] Mitchell C, Pritchard-Jones K, Shannon R, Hutton C, Stevens S, Machin D, Imeson J, Kelsey A, Vujanic GM, Gornall P, Walker J, Taylor R, Sartori P, Hale J, Levitt G, Messahel B, Middleton H, Grundy R, Pritchard J; United Kingdom Cancer Study Group. Immediate nephrectomy versus preoperative chemotherapy in the management of non-metastatic Wilms tumour: results of a randomised trial (UKW3) by the UK Children's Cancer Study Group. *Eur J Cancer.* 2006 Oct; 42(15): 2554-62.

[17] Somers JM, Lomas DJ, Hacking JC, Coleman N, Broadbent VA, Dixon AK. Radiologically-guided cutting needle biopsy for suspected malignancy in childhood. *Clin Radiol.* 1993 Oct; 48(4): 236-40.

[18] UMBRELLA Protocol SIOP-RTSG 2016, Integrated research and guidelines for standardized diagnostics and therapy.{<http://www.fnkc.ru/docs/SIOP-RTSG2016.pdf>} (Accessed on 12 February 2024).

**Address for communication:** Dr Yogesh Kumar Sarin,  
Email: [yksarin@gmail.com](mailto:yksarin@gmail.com)

© Authors; Distributed under Creative Commons CC-BY-NC-ND attribution 4.0 international license

Received 2 March 2024; Accepted 12 March 2024

Acknowledgements : None.

Conflicts of Interest : None declared by authors

Source of Funding : None

Ethical concerns : Institute Ethical Committee approval number LHMCI/EC/2021/03/88

**Citation:** Singh P, Sarin YK, Nangia A. A study of retro-peritoneal core needle biopsy in the diagnosis of suspected malignant renal tumors. *Pediatr Surg Trop* 2024; 1(2): 102-107.

